

Somatic Accumulation of GluA1-AMPA Receptors Leads to Selective Cognitive Impairments in Mice

Bannerman D., Borchardt T., Jensen V., Rozov A., Haj-Yasein N., Burnashev N., Zamanillo D., Bus T., Grube I., Adelmann G., Rawlins J., Sprengel R.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Bannerman, Borchardt, Jensen, Rozov, Haj-Yasein, Burnashev, Zamanillo, Bus, Grube, Adelmann, Rawlins and Sprengel. The GluA1 subunit of the L- α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor (AMPA) plays a crucial, but highly selective, role in cognitive function. Here we analyzed AMPAR expression, AMPAR distribution and spatial learning in mice (Gria1R/R), expressing the “trafficking compromised” GluA1(Q600R) point mutation. Our analysis revealed somatic accumulation and reduction of GluA1(Q600R) and GluA2, but only slightly reduced CA1 synaptic localization in hippocampi of adult Gria1R/R mice. These immunohistological changes were accompanied by a strong reduction of somatic AMPAR currents in CA1, and a reduction of plasticity (short-term and long-term potentiation, STP and LTP, respectively) in the CA1 subfield following tetanic and theta-burst stimulation. Nevertheless, spatial reference memory acquisition in the Morris water-maze and on an appetitive Y-maze task was unaffected in Gria1R/R mice. In contrast, spatial working/short-term memory during both spontaneous and rewarded alternation tasks was dramatically impaired. These findings identify the GluA1(Q600R) mutation as a loss of function mutation that provides independent evidence for the selective role of GluA1 in the expression of short-term memory.

<http://dx.doi.org/10.3389/fnmol.2018.00199>

Keywords

AMPA receptors, GluA1, Long-term potentiation, Morris water-maze, RNA-editing, Spatial memory, Spatial working memory

References

- [1] Andrásfalvy B. K., Smith M. A., Borchardt T., Sprengel R., Magee J. C., (2003). Impaired regulation of synaptic strength in hippocampal neurons from GluR1-deficient mice. *J. Physiol.* 552, 35–45. 10.1113/jphysiol.2003.04557512878757.
- [2] Bannerman D. M., Deacon R. M., Brady S., Bruce A., Sprengel R., Seeburg P. H., (2004). A comparison of GluR1-deficient and wild-type mice on a test battery assessing sensorimotor, affective, and cognitive behaviors. *Behav. Neurosci.* 118, 643–647. 10.1037/0735-7044.118.3.64315174943.
- [3] Bliss T. V., Lomo T., (1973). Long-lasting potentiation of synaptic transmission in the dentate area of the anaesthetized rabbit following stimulation of the perforant path. *J. Physiol.* 232, 331–356. 10.1113/jphysiol.1973.sp0102734727084.

- [4] Brechet A., Buchert R., Schwenk J., Boudkazi S., Zolles G., Siquier-Pernet K.. (2017). AMPA-receptor specific biogenesis complexes control synaptic transmission and intellectual ability. *Nat. Commun.* 8:15910. 10.1038/ncomms1591028675162.
- [5] Brusa R., Zimmermann F., Koh D. S., Feldmeyer D., Gass P., Seeburg P. H.. (1995). Early-onset epilepsy and postnatal lethality associated with an editing-deficient GluR-B allele in mice. *Science* 270, 1677-1680. 10.1126/science.270.5242.16777502080.
- [6] Burnashev N., Monyer H., Seeburg P. H., Sakmann B., (1992). Divalent ion permeability of AMPA receptor channels is dominated by the edited form of a single subunit. *Neuron* 8, 189-198. 10.1016/0896-6273(92)90120-31370372.
- [7] Deacon R. M., Rawlins J. N., (2006). T-maze alternation in the rodent. *Nat. Protoc.* 1, 7-12. 10.1038/nprot.2006.217406205.
- [8] Fuchs E. C., Zivkovic A. R., Cunningham M. O., Middleton S., Lebeau F. E., Bannerman D. M.. (2007). Recruitment of parvalbumin-positive interneurons determines hippocampal function and associated behavior. *Neuron* 53, 591-604. 10.1016/j.neuron.2007.01.03117296559.
- [9] Gerlai R., Henderson J. T., Roder J. C., Jia Z., (1998). Multiple behavioral anomalies in GluR2 mutant mice exhibiting enhanced LTP. *Behav. Brain Res.* 95, 37-45. 10.1016/s0166-4328(98)00002-39754875.
- [10] Granger A. J., Shi Y., Lu W., Cerpas M., Nicoll R. A., (2013). LTP requires a reserve pool of glutamate receptors independent of subunit type. *Nature* 493, 495-500. 10.1038/nature1177523235828.
- [11] Greger I. H., Khatri L., Kong X., Ziff E. B., (2003). AMPA receptor tetramerization is mediated by Q/R editing. *Neuron* 40, 763-774. 10.1016/s0896-6273(03)00668-814622580.
- [12] Greger I. H., Khatri L., Ziff E. B., (2002). RNA editing at arg607 controls AMPA receptor exit from the endoplasmic reticulum. *Neuron* 34, 759-772. 10.1016/s0896-6273(02)00693-112062022.
- [13] Higuchi M., Maas S., Single F. N., Hartner J., Rozov A., Burnashev N.. (2000). Point mutation in an AMPA receptor gene rescues lethality in mice deficient in the RNA-editing enzyme ADAR2. *Nature* 406, 78-81. 10.1038/3501755810894545.
- [14] Hoffman D. A., Sprengel R., Sakmann B., (2002). Molecular dissection of hippocampal theta-burst pairing potentiation. *Proc. Natl. Acad. Sci. U S A* 99, 7740-7745. 10.1073/pnas.09215799912032353.
- [15] Jensen V., Kaiser K. M., Borchardt T., Adelmann G., Rozov A., Burnashev N.. (2003). A juvenile form of postsynaptic hippocampal long-term potentiation in mice deficient for the AMPA receptor subunit GluR-A. *J. Physiol.* 553, 843-856. 10.1113/jphysiol.2003.05363714555717.
- [16] Jia Z., Agopyan N., Miu P., Xiong Z., Henderson J., Gerlai R.. (1996). Enhanced LTP in mice deficient in the AMPA receptor GluR2. *Neuron* 17, 945-956. 10.1016/s0896-6273(00)80225-18938126.
- [17] Kask K., Zamanillo D., Rozov A., Burnashev N., Sprengel R., Seeburg P. H., (1998). The AMPA receptor subunit GluR-B in its Q/R site-unedited form is not essential for brain development and function. *Proc. Natl. Acad. Sci. U S A* 95, 13777-13782. 10.1073/pnas.95.23.137779811877.
- [18] Keinänen K., Wisden W., Sommer B., Werner P., Herb A., Verdoorn T. A.. (1990). A family of AMPA-selective glutamate receptors. *Science* 249, 556-560. 10.1126/science.21663372166337.
- [19] Larson J., Lynch G., (1986). Induction of synaptic potentiation in hippocampus by patterned stimulation involves two events. *Science* 232, 985-988. 10.1126/science.37046353704635.
- [20] Lu W., Shi Y., Jackson A. C., Bjorgan K., During M. J., Sprengel R.. (2009). Subunit composition of synaptic AMPA receptors revealed by a single-cell genetic approach. *Neuron* 62, 254-268. 10.1016/j.neuron.2009.02.02719409270.
- [21] Luchkina N. V., Coleman S. K., Huupponen J., Cai C., Kivistö A., Taira T.. (2017). Molecular mechanisms controlling synaptic recruitment of GluA4 subunit-containing AMPA-receptors critical for functional maturation of CA1 glutamatergic synapses. *Neuropharmacology* 112, 46-56. 10.1016/j.neuropharm.2016.04.04927157711.
- [22] Madeo M., Stewart M., Sun Y., Sahir N., Wiethoff S., Chandrasekar I.. (2016). Loss-of-function mutations in FRRS1L lead to an epileptic-dyskinetic encephalopathy. *Am. J. Hum. Genet.* 98, 1249-1255. 10.1016/j.ajhg.2016.04.00827236917.
- [23] Monyer H., Seeburg P. H., Wisden W., (1991). Glutamate-operated channels: developmentally early and mature forms arise by alternative splicing. *Neuron* 6, 799-810. 10.1016/0896-6273(91)90176-z1673851.
- [24] Morris R. G. M., (1981). Spatial localization does not require the presence of local cues. *Learn. Motiv.* 12, 239-260. 10.1016/0023-9690(81)90020-5.
- [25] Morris R. G., Garrud P., Rawlins J. N., O'Keefe J., (1982). Place navigation impaired in rats with hippocampal lesions. *Nature* 297, 681-683. 10.1038/297681a07088155.
- [26] Nagy A., Rossant J., Nagy R., Abramow-Newerly W., Roder J. C., (1993). Derivation of completely cell culture-derived mice from early-passage embryonic stem cells. *Proc. Natl. Acad. Sci. U S A* 90, 8424-8428. 10.1073/pnas.90.18.84248378314.

- [27] Pelkey K. A., Barksdale E., Craig M. T., Yuan X., Sukumaran M., Vargish G. A.. (2016). Pentraxins coordinate excitatory synapse maturation and circuit integration of parvalbumin interneurons. *Neuron* 90:661. 10.1016/j.neuron.2016.04.02427151642.
- [28] Plant K., Pelkey K. A., Bortolotto Z. A., Morita D., Terashima A., McBain C. J.. (2006). Transient incorporation of native GluR2-lacking AMPA receptors during hippocampal long-term potentiation. *Nat. Neurosci.* 9, 602-604. 10.1038/nn167816582904.
- [29] Reisel D., Bannerman D. M., Schmitt W. B., Deacon R. M., Flint J., Borchardt T.. (2002). Spatial memory dissociations in mice lacking GluR1. *Nat. Neurosci.* 5, 868-873. 10.1038/nn91012195431.
- [30] Romberg C., Raffel J., Martin L., Sprengel R., Seeburg P. H., Rawlins J. N.. (2009). Induction and expression of GluA1 (GluR-A)-independent LTP in the hippocampus. *Eur. J. Neurosci.* 29, 1141-1152. 10.1111/j.1460-9568.2009.06677.x19302150.
- [31] Rozov A., Sprengel R., Seeburg P. H., (2012). GluA2-lacking AMPA receptors in hippocampal CA1 cell synapses: evidence from gene-targeted mice. *Front. Mol. Neurosci.* 5:22. 10.3389/fnmol.2012.0002222375105.
- [32] Sanderson D. J., Good M. A., Skelton K., Sprengel R., Seeburg P. H., Rawlins J. N.. (2009). Enhanced long-term and impaired short-term spatial memory in GluA1 AMPA receptor subunit knockout mice: evidence for a dual-process memory model. *Learn. Mem.* 16, 379-386. 10.1101/lm.133910919470654.
- [33] Sanderson D. J., Gray A., Simon A., Taylor A. M., Deacon R. M., Seeburg P. H.. (2007). Deletion of glutamate receptor-A (GluR-A) AMPA receptor subunits impairs one-trial spatial memory. *Behav. Neurosci.* 121, 559-569. 10.1037/0735-7044.121.3.55917592947.
- [34] Sanderson D. J., Hindley E., Smeaton E., Denny N., Taylor A., Barkus C.. (2011a). Deletion of the GluA1 AMPA receptor subunit impairs recency-dependent object recognition memory. *Learn. Mem.* 18, 181-190. 10.1101/lm.208341121378100.
- [35] Sanderson D. J., Sprengel R., Seeburg P. H., Bannerman D. M., (2011b). Deletion of the GluA1 AMPA receptor subunit alters the expression of short-term memory. *Learn. Mem.* 18, 128-131. 10.1101/lm.201491121325433.
- [36] Sanderson D. J., McHugh S. B., Good M. A., Sprengel R., Seeburg P. H., Rawlins J. N.. (2010). Spatial working memory deficits in GluA1 AMPA receptor subunit knockout mice reflect impaired short-term habituation: evidence for Wagner's dual-process memory model. *Neuropsychologia* 48, 2303-2315. 10.1016/j.neuropsychologia.2010.03.01820350557.
- [37] Schmitt W. B., Deacon R. M., Seeburg P. H., Rawlins J. N., Bannerman D. M., (2003). A within-subjects, within-task demonstration of intact spatial reference memory and impaired spatial working memory in glutamate receptor-A-deficient mice. *J. Neurosci.* 23, 3953-3959. 10.1523/JNEUROSCI.23-09-03953.200312736365.
- [38] Sommer B., Köhler M., Sprengel R., Seeburg P. H., (1991). RNA editing in brain controls a determinant of ion flow in glutamate-gated channels. *Cell* 67, 11-19. 10.1016/0092-8674(91)90568-j1717158.
- [39] Soto D., Coombs I. D., Kelly L., Farrant M., Cull-Candy S. G., (2007). Stargazin attenuates intracellular polyamine block of calcium-permeable AMPA receptors. *Nat Neurosci* 10, 1260-1267. 10.1038/nn196617873873.
- [40] Soto D., Coombs I. D., Renzi M., Zonouzi M., Farrant M., Cull-Candy S. G., (2009). Selective regulation of long-form calcium-permeable AMPA receptors by an atypical TARP, γ -5. *Nat. Neurosci.* 12, 277-285. 10.1038/nn.226619234459.
- [41] Sprengel R., Aronoff R., Völkner M., Schmitt B., Mosbach R., Kuner T., (2001). Glutamate receptor channel signatures. *Trends Pharmacol. Sci.* 22, 7-10. 10.1016/s0165-6147(00)01588-111165660.
- [42] Swanson G. T., Kamboj S. K., Cull-Candy S. G., (1997). Single-channel properties of recombinant AMPA receptors depend on RNA editing, splice variation and subunit composition. *J. Neurosci.* 17, 58-69. 10.1523/JNEUROSCI.17-01-00058.19978987736.
- [43] Vekovischeva O. Y., Aitta-Aho T., Echenko O., Kankaanpää A., Seppälä T., Honkanen A.. (2004). Reduced aggression in AMPA-type glutamate receptor GluR-A subunit-deficient mice. *Genes Brain Behav.* 3, 253-265. 10.1111/j.1601-1848.2004.00075.x15344919.
- [44] Vekovischeva O. Y., Zamanillo D., Echenko O., Seppälä T., Uusi-Oukari M., Honkanen A.. (2001). Morphine-induced dependence and sensitization are altered in mice deficient in AMPA-type glutamate receptor-A subunits. *J. Neurosci.* 21, 4451-4459. 10.1523/JNEUROSCI.21-12-04451.200111404432.
- [45] Wenthold R. J., Petralia R. S., Blahos J., II, Niedzielski A. S., (1996). Evidence for multiple AMPA receptor complexes in hippocampal CA1/CA2 neurons. *J. Neurosci.* 16, 1982-1989. 10.1523/JNEUROSCI.16-0-01982.19968604042.
- [46] Wigström H., Gustafsson B., (1983). Facilitated induction of hippocampal long-lasting potentiation during blockade of inhibition. *Nature* 301, 603-604. 10.1038/301603a06298626.
- [47] Wigström H., Gustafsson B., (1985). Facilitation of hippocampal long-lasting potentiation by GABA antagonists. *Acta Physiol. Scand.* 125, 159-172. 10.1111/j.1748-1716.1985.tb07703.x2996303.
- [48] Zamanillo D., Sprengel R., Hvalby O., Jensen V., Burnashev N., Rozov A.. (1999). Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* 284, 1805-1811. 10.1126/science.284.5421.180510364547.

- [49] Zhu J. J., Esteban J. A., Hayashi Y., Malinow R., (2000). Postnatal synaptic potentiation: delivery of GluR4-containing AMPA receptors by spontaneous activity. *Nat. Neurosci.* 3, 1098-1106. 10.1038/8061411036266.